

Multiple Sclerosis Bibliography

- Armutlu, K., Karabudak, R., & Nurlu, G. (2001). Physiotherapy approaches in the treatment of ataxic Multiple Sclerosis: A pilot study. *NeuroRehabilitation and Neural Repair, 15*(3), 203-211.
- Brown, T., & Kraft, G. (2005). Exercise and rehabilitation for individuals with Multiple Sclerosis. *Phys Med Rehabil Clin N Am, 16*, 513-555.
- Cattaneo, D., Jonsdittor, J., Zocchi, M., & Regola, A. (2007). Effects of balance exercises on people with Multiple Sclerosis: A pilot study. *Clinical Rehabilitation, 21*, 771-781.
- Cattaneo, D., Jonsdittor, J., & Repetti, S. (2007). Reliability of four scales on balance disorders in persons with Multiple Sclerosis. *Disability and Rehabilitation, 29*(24), 1920-1925.
- Cattaneo, D., Regola, A., & Meotti, M. (2006). Validity of six balance disorders scales in persons with Multiple Sclerosis. *Disability and Rehabilitation, 28*(12), 789-795.
- DeBolt, L., & McCubbin, J. (2004). The effects of home-based resistance exercise on balance, power, and mobility in adults with Multiple Sclerosis. *Arch Phys Med Rehabil, 85*, 290-297.
- Finlayson, M., Peterson, E., & Cho, C. (2006). Risk factors for falling among people aged 45 to 90 years with Multiple Sclerosis. *Arch Phys Med Rehabil, 87*, 1274-1279.
- Fulk, G. (2005). Locomotor training and virtual reality-based balance training for an individual with Multiple Sclerosis: A case report. *Journal of Neurological Physical Therapy, 29*(1), 34-42.

- McConvey, J., & Bennet, S. (2005). Reliability of the dynamic gait index in individuals with Multiple Sclerosis. *Arch Phys Med Rehabil*, 86, 130-133.
- Nieuwenhuis, M., Van Tongeren, H., Sorensen, P., & Raunborg, M. (2006). The six spot step test: A new measurement for walking ability in Multiple Sclerosis. *Multiple Sclerosis*, 12, 495-500.
- Paltamaa, J., Sarasoja, T., Leskinen, E., Wikstrom, J., & Malkia, E. (2007). Measures of physical functioning predict self-reported performance in self care, mobility, and domestic life in ambulatory persons with Multiple Sclerosis. *Arch Phys Med Rehabil*, 88, 1649-1657.
- Peterson, E., Cho, C., & Finlayson, M. (2007). Fear of falling and associated activity curtailment among middle aged and older adults with Multiple Sclerosis. *Multiple Sclerosis*, 13, 1168-1175.
- Romberg, A., Virtanen, A., Ruutiainen, J., Aunola, S., Karppi, S., Vaara, M., Surakka, J., Pohjolainen, T., & Seppanen, A. (2004). Effects of a 6-month exercise program on patients with Multiple Sclerosis. *Neurology*, 63, 2034-2038.
- Silkwood-Sherer, D., & Warmbier, H. (2007). Effects of hippotherapy on postural stability in persons with Multiple Sclerosis: A pilot study. *JNPT*, 31, 77-84.
- Soyuer, F., Mirza, M., & Erkorkmaz, U. (2006). Balance performance in three forms of Multiple Sclerosis. *Neurological Research*, 28, 555-562.
- White, L., & Dressendorfer, R. (2004). Exercise and Multiple Sclerosis. *Sports Medicine*, 34(15), 1077-1100.

